



Vegetation Continuous Fields


Evolution through multiple spatial and temporal resolutions




John Townshend¹, Charlene DiMiceli¹, Robert Sohlberg¹, Mark Carroll², Do-Hyung Kim¹, Chengquan Huang¹, Matthew Hansen¹

¹ University of Maryland, Department of Geographical Sciences, ² Sigma Space, NASA Goddard Space Flight Center


AVHRR VCF







MODIS VCF



Landsat VCF



Sexton et al. 2013







Algorithm / Resolution
Linear Mixture Model / 1 km

Regression Tree / 500 m

Bagged Regression Tree
(Fully Automated) / 250 m

Regression Tree (trained with
250 m MODIS) / 30 m

High Res. Reference




World imagery (ESRI, 2014)

30 years (1982 – 2014),
annual, 1/20 Degree

A time series of MODIS VCF V5 yearly data is used to identify deforested areas (X.-P. Song et al., in preparation)

Applications



Details of VCFs

Instrument	Spatial Resolution	Spatial & Temporal Extent	References
AVHRR (GVI prototype)	1°	North America subset, 1989	DeFries et al. 1995
AVHRR (Pathfinder prototype)	8 km	Central Africa, 1987	DeFries et al. 1997
AVHRR	8 km	Global, 1982 – 1994	DeFries et al. 2000
AVHRR	1 km	Global, 1992 – 1993	DeFries et al. 2000
MODIS	500 m	Global, 2001	Hansen et al. 2003
MODIS	250 m	Global, annual 2000-2010	DiMiceli, et al. 2011
AVHRR & MODIS	1/20°	Global, annual, 1982-2014	In Preparation
LANDSAT	30 m	Global, 2000, 2005	Sexton, et al. 2013 http://www.landcover.org
LANDSAT	30 m	Global 2000	Hansen, et al. 2013 http://earthenginepartners.appspot.com/science-2013-global-forest

Acknowledgement

This study is funded under NASA grants NNX11AE77G, NNX13AS4SA , NNH06ZDA001N